

# Management of Diabetes Undergoing Elective Surgery

## Diabetes and Endocrinology

Patient Information Leaflet for patients on basal/bolus insulin (≥ 4 injections per day)

These instructions are for patients with diabetes, treated with both:

 Bolus insulin for mealtimes (eg Novorapid, Humalog, Apidra)

#### And

 Background insulin (eg Lantus, Levemir, Insulatard, Humulin I, Degludec, Toujeo).

#### On the day before surgery

If you are on dapaglifozin, empagliflozin or canagliflozin, stop taking these drugs until after your operation and normal diet resumes.

Continue your usual diabetic medications and follow any bowel preparations as advised by your surgeon / staff at the pre-assessment clinic.

### On the day of surgery

When your background insulin is due, you should take these as normal.

You should **not** take your mealtime bolus insulin from the time you are asked to fast.

If you are on dapagliflozin, canagliflozin and empagliflozin, you would have stopped taking these the day before.

Under special circumstances, your metformin may need to be withheld for 48 hours. Staff from the pre-assessment team will inform you of this.

You can continue taking other diabetic medication as usual.

Remember to bring to hospital all of your diabetes medications, dextrose tablets and blood testing equipment.

Your blood glucose will be tested by finger prick when you arrive to hospital and then monitored.

You can ask for extra blood testing if you feel symptoms of a 'hypo' (low blood glucose).

The target for safe surgery is 6-10 mmol. You may need insulin treatment to bring down high glucose, or dextrose infusion for treatment of low blood glucose.

#### **After surgery**

You will be discharged when you are eating and drinking normally and your blood glucose is at a safe level.

However, if you are unwell with:

- Continuous vomiting / diarrhoea, or high fever
- Unable to keep food down for four hours or more
- **High** blood glucose (>15.0 mmol by finger prick testing)
- **High** ketones (>0.6 mmol)

You should seek medical help. During office hours, contact your usual diabetes specialist nurse or doctor.

At other times, contact your local out-of-hours GP service.

If you have any questions, or if there is anything you do not understand about this leaflet, please contact:

The Diabetes and Endocrine Centre on 01384 244399

Russells Hall Hospital switchboard number: 01384 456111

#### This leaflet can be downloaded or printed from:

http://dgft.nhs.uk/services-and-wards/diabetes-and-endocrinology/

If you have any feedback on this patient information leaflet, please email dgft.patient.information@nhs.net

This leaflet can be made available in large print, audio version and in other languages, please call 0800 073 0510.

للحصول على هذه النشرة بحجم أكبر، وعلى شكل إصدار صوتي و بلغات أخرى، الرجاء الاتصال بالرقم 08000730510.

此宣传单可提供大字版本、音频版本和其它语言版本,请拨打电话: 0800 073 0510。

Ulotka dostępna jest również w dużym druku, wersji audio lub w innym języku. W tym celu zadzwoń pod numer 0800 073 0510.

ਇਹ ਪਰਚਾ ਵੱਡੇ ਅੱਖਰਾਂ, ਬੋਲ ਕੇ ਰੀਕਾਰਡ ਕੀਤਾ ਹੋਇਆ ਅਤੇ ਦੂਸਰੀਆਂ ਭਾਸ਼ਾਵਾਂ ਵਿਚ ਵੀ ਪਾਪਤ ਹੋ ਸਕਦਾ ਹੈ, 0800 073 0510 ਤੇ ਫੋਨ ਕਰੋ ਜੀ।

Aceasta brosura poate fi pusa la dispozitie tiparita cu caractere mari, versiune audio sau in alte limbi, pentru acest lucru va rugam sunati la 0800 073 0510.

یہ کتابچہ آپ کو بڑے حروف کی لکھائی ، سمعی صورت اور دیگر زبانوں میں مہیا کیا جا سکتا ہے برائے مہربانی فون نمبر 08000730510 پر رابطہ کریں۔

Originator: Terence Pang, Chris Homer. Date originated: October 2013. Date reviewed: June 2021. Next review date: March 2024. Version: 5. DGH ref: DGH/PIL/00897.