

Thrombolysis

Radiology Department Patient Information Leaflet

Introduction

This leaflet tells you about having thrombolysis. It explains what is involved and what the possible risks are. It is not meant to replace informed discussion between you and your doctor, but can act as a starting point for such discussions. If you have any questions about the procedure, please ask the doctor who has referred you or the department which is going to perform it.

What is thrombolysis?

Thrombolysis is the breakdown of blood clots with the use of drugs. Once a clot starts to form in a blood vessel, it may continue and block the whole vessel. While an operation may be necessary to remove the clot, it is also possible to dissolve the clot by directly injecting a special 'clot-busting' drug into the artery or vein. This can lead to a great improvement in blood flow and may avoid the need for an operation. Sometimes, an underlying narrowing is revealed in the blood vessel once the clot has dissolved, and it may be possible to treat this by angioplasty (balloon) or stent insertion at the same time.

Why do you need thrombolysis?

Your doctors know that there is a blockage in a blood vessel based on the tests you have already had. These tests might include a Doppler ultrasound scan, magnetic resonance imaging (MRI) or computed tomography (CT) angiography. If nothing is done about the situation, then severe and permanent damage may occur. While the blockage could need treatment with surgery, in your case it has been decided that thrombolysis is the best way of proceeding.

Are there any risks?

Thrombolysis is generally a safe procedure, but as with any medical procedure, there are some risks and complications that can arise. Bruising at the puncture site is very common and rarely may require a small operation. Occasionally, ongoing leakage from the puncture site may cause a swelling called a 'false aneurysm' that may require a further procedure.

Clot-busting drugs have to be very powerful to work, consequently there is a risk that bleeding will occur elsewhere in your body. Commonly, this is from the bowel and might require treatment or stopping the thrombolysis. Very rarely, bleeding can occur in the brain and cause a stroke. There may be a risk to life, but again, this is very rare.

Sometimes, the blood clot may be so extensive that the clot-busting drug simply cannot dissolve it all away. In these cases, surgery may be required to relieve the blockage.

If angioplasty or venoplasty is required, there are additional risks related to these procedures (see the relevant patient information leaflets).

Who has made the decision?

The consultant in charge of your care and the interventional radiologist performing the procedure have discussed your case and feel that this is the best option.

However, you will also have the opportunity for your opinion to be considered and if, after discussion with your doctors, you no longer want the procedure, you can decide against it.

Are you required to make any special preparations?

Thrombolysis is usually carried out under local anaesthetic. You may be asked not to eat for four hours before the procedure, although you may still drink clear fluids, such as water.

If you have any allergies or have previously had a reaction to the dye (contrast agent), you must tell the radiology staff before you have the test.

Who will you see?

You will be seen by a specially trained team led by an interventional radiologist within the radiology department. Interventional radiologists have special expertise in reading the images and using imaging to guide catheters and wires to aid diagnosis and treatment.

Where will the procedure take place?

The procedure will take place in the angiography suite or theatre. This is usually located within the radiology department. This is similar to an operating theatre, into which specialised X-ray equipment has been installed.

What happens during thrombolysis?

Before the procedure, the interventional radiologist will explain the procedure and ask you to sign a consent form. Please feel free to ask any questions that you may have and remember that even at this stage, you can decide against going ahead with the procedure if you so wish.

You will be asked to get undressed and put on a hospital gown. A small cannula (thin tube) may be placed into a vein in your arm. You may receive a sedative to relieve anxiety, as well as an antibiotic. You will lie on the X-ray table, generally flat on your back. It is normal to remain awake during the procedure. Your pulse, blood pressure, electrocardiogram (ECG), and blood oxygen levels will be monitored throughout the procedure and you may be given oxygen.

The procedure is performed under sterile conditions and the interventional radiologist and radiology nurse will wear sterile gowns and gloves to carry out the procedure. The skin near the point of insertion, usually the groin, will be swabbed with antiseptic and you will be covered with sterile drapes. The skin and deeper tissues over the artery or vein will be numbed with local anaesthetic, and a fine plastic tube (catheter) is then passed over a wire and into the artery or vein.

The radiologist will use the X-ray equipment and small amounts of dye (contrast agent) to make sure that the catheter is moved into the right position, very close or actually into the blood clot. The clot-busting drug (thrombolytic) is injected down the catheter and into the blood clot. The catheter is left in the artery or vein and attached to an infusion pump, so that injection of the clot-busting drug can be continued over hours (occasionally up to 48 to 72 hours), during which time you will be transferred back to the ward for careful observation.

The radiologist will check progress periodically by injecting the dye to show how much of the clot has dissolved. Once the procedure is completed, the catheter will be removed and firm pressure will be applied for about 10 minutes to prevent any bleeding.

Will it hurt?

When the local anaesthetic is injected, it will sting for a short while, but this soon wears off. You may feel a warm sensation for a few seconds when the dye is injected, and it may feel like you are passing urine.

How long will it take?

Every patient is different, and it is not always easy to predict. However, expect to be in the radiology department for at least an hour initially. You may require several trips to the X-ray department to check on progress. Sometimes, an underlying narrowing is revealed once the clot has cleared, which may be treated at the time by angioplasty or stent.

X-ray precautions

The procedure uses X-rays, and these use a small amount of radiation which may add slightly to the normal risk of cancer.

Your doctor thinks that the benefit of the examination outweighs the risks. However, if you are concerned about these possible risks, please discuss these with this doctor.

For more information, visit:

<https://www.gov.uk/government/publications/medical-radiation-patient-doses/patient-dose-information-guidance>

If you are a female patient, you must tell us if you are or might be pregnant before you attend for your scan.

Finally

Some of your questions should have been answered by this leaflet, but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. Make sure you are satisfied that you have received enough information about the procedure.

Contact

The British Society of Interventional Radiology

www.bsir.org

Department of Radiology
Russells Hall Hospital
West Wing
Ground Floor
Pensnett Road
Dudley
West Midlands
DY1 2HQ

© This leaflet has been reproduced with kind permission from British
Society of Interventional Radiology

If you have any questions, or if there is anything you do not understand about this leaflet, please contact:

The Radiology Department

Russells Hall Hospital switchboard number: 01384 456111

This leaflet can be downloaded or printed from:

<http://dgft.nhs.uk/services-and-wards/radiology>

If you have any feedback on this patient information leaflet, please email dgft.patient.information@nhs.net

This leaflet can be made available in large print, audio version and in other languages, please call 0800 073 0510.

للحصول على هذه النشرة بحجم أكبر، وعلى شكل إصدار صوتي و بلغات أخرى، الرجاء الاتصال بالرقم 08000730510.

此宣传单可提供大字版本、音频版本和其它语言版本，请拨打电话：0800 073 0510。

Ulotka dostępna jest również w dużym druku, wersji audio lub w innym języku. W tym celu zadzwoń pod numer 0800 073 0510.

ਇਹ ਪਰਚਾ ਵੱਡੇ ਅੱਖਰਾਂ, ਬੋਲ ਕੇ ਰੀਕਾਰਡ ਕੀਤਾ ਹੋਇਆ ਅਤੇ ਦੂਸਰੀਆਂ ਭਾਸ਼ਾਵਾਂ ਵਿਚ ਵੀ ਪ੍ਰਾਪਤ ਹੋ ਸਕਦਾ ਹੈ, 0800 073 0510 ਤੇ ਫੋਨ ਕਰੋ ਜੀ।

Aceasta broșura poate fi pusă la dispoziție tipărită cu caractere mari, versiune audio sau în alte limbi, pentru acest lucru vă rugăm sunați la 0800 073 0510.

یہ کتابچہ آپ کو بڑے حروف کی لکھائی، سمعی صورت اور دیگر زبانوں میں مہیا کیا جا سکتا ہے۔ براہ کرم 08000730510 پر رابطہ کریں۔