

# Transjugular Liver Biopsy

## Radiology Department Patient Information Leaflet

### Introduction

This leaflet tells you about having a transjugular liver biopsy. It explains what is involved and what the possible risks are. It is not meant to replace informed discussion between you and your doctor, but can act as a starting point for such discussions. If you have any questions about the procedure, please ask the doctor who has referred you or the department that will be performing it.

### What is a transjugular liver biopsy?

A liver biopsy is a procedure that involves taking a tiny specimen of the liver for examination under a microscope. In most cases, a liver biopsy is taken through the skin by passing a fine needle through into the liver.

A transjugular liver biopsy is an alternative way of obtaining the liver specimen by passing the needle through the vein in the neck (jugular vein). This method is used in patients who have abnormal clotting of the blood or fluid collecting within the abdomen. This technique is used to reduce the risk of bleeding after the biopsy.

## Why do you need a transjugular liver biopsy?

The doctors looking after you have decided that you need a liver biopsy to obtain more information about your liver problem. The information gained from the biopsy will help the doctors in the treatment of your condition.

## Are there any risks?

Transjugular liver biopsy is a safe procedure, but as with any medical procedure, there are some risks and complications that can arise. The overall risk of a problem requiring further treatment is low (one to two per cent). The main risk is bleeding after the biopsy, and rarely this can cause a risk to life. However, a transjugular liver biopsy has a lower risk of bleeding than a conventional liver biopsy taken through the side of the abdomen.

## Who has made the decision?

The consultant in charge of your care and the interventional radiologist performing the procedure have discussed your case and feel that this is the best option. However, you will also have the opportunity for your opinion to be considered and if, after discussion with your doctors, you do not want the procedure carried out, you can decide against it.

## Are you required to make any special preparations?

A transjugular liver biopsy is usually carried out as a day case procedure under local anaesthetic. You may be asked not to eat for four hours before the procedure, although you may still drink clear fluids, such as water. You may receive a sedative to relieve anxiety, as well as an antibiotic.

If your blood clotting is abnormal, you may be given special blood transfusions to try and correct this. If you have any concerns about having blood transfusions, you should discuss these with your doctor. If you have any allergies or have previously had a reaction to the dye (contrast agent), you **must** tell the radiology staff before you have the test.

## Who will you see?

You will see a specially trained team led by an interventional radiologist within the radiology department. Interventional radiologists have special expertise in reading the images and using imaging to guide catheters and wires to aid diagnosis and treatment.

## Where will the procedure take place?

The procedure will take place in the interventional theatre, located within the radiology department. This is similar to an operating theatre, into which specialised X-ray equipment has been installed.

## What happens during a transjugular liver biopsy?

You will be asked to get undressed and put on a hospital gown. A small cannula (thin tube) may be placed into a vein in your arm. The procedure is performed using local anaesthetic. The skin at the side of the neck will be swabbed and covered with sterile towels. The skin will be numbed with local anaesthetic.

Once the skin is numb, a small tube (catheter) is inserted into the vein at the side of the neck. An X-ray machine is used to guide the catheter into the vein in the liver, and then to guide the needle into the liver. Usually, two to three biopsy specimens are taken.

## Will it hurt?

When the local anaesthetic is injected, it will sting for a short while, but this soon wears off. When the catheter is placed in the liver, you may get a dull ache in your right shoulder. This will go away once the tube has been removed. Some people feel a sharp pinch inside the abdomen as the biopsy is taken, but this will only last for one to two seconds.

## How long will it take?

Every patient is different, and it is not always easy to predict; however, expect to be in the radiology department for about one hour altogether.

## What happens afterwards?

You will be taken back to your ward on a trolley. Nurses on the ward will carry out routine observations, such as taking your pulse and blood pressure, to make sure that there are no problems. You will generally stay in bed for a few hours, until you have recovered. Assuming you are feeling well, you will normally be discharged the same day.

## How will I get my results?

The biopsy specimens will be sent for examination. Once the results are available, your doctors will discuss these with you.

## X-ray precautions

The procedure uses X-rays, and these use a small amount of radiation which may add slightly to the normal risk of cancer.

Your doctor thinks that the benefit of the examination outweighs the risks. However, if you are concerned about these possible risks, please discuss these with this doctor.

For more information visit:

<https://www.gov.uk/government/publications/medical-radiation-patient-doses/patient-dose-information-guidance>

**If you are a female patient, you must tell us if you are or might be pregnant before you attend for your scan.**

## Finally

Some of your questions should have been answered by this leaflet, but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. Make sure you are satisfied that you have received enough information about the procedure.

## Contact

The British Society of Interventional Radiology

[www.bsir.org](http://www.bsir.org)

Department of Radiology  
Russells Hall Hospital  
West Wing  
Ground Floor  
Pensnett Road  
Dudley  
West Midlands  
DY1 2HQ

© This leaflet has been reproduced with kind permission from British  
Society of Interventional Radiology

If you have any questions, or if there is anything you do not understand about this leaflet, please contact:

The Radiology Department

Russells Hall Hospital switchboard number: 01384 456111

**This leaflet can be downloaded or printed from:**

<http://dgft.nhs.uk/services-and-wards/radiology>

If you have any feedback on this patient information leaflet, please email [dgft.patient.information@nhs.net](mailto:dgft.patient.information@nhs.net)

**This leaflet can be made available in large print, audio version and in other languages, please call 0800 073 0510.**

للحصول على هذه النشرة بحجم أكبر، وعلى شكل إصدار صوتي و بلغات أخرى، الرجاء الاتصال بالرقم 08000730510.

此宣传单可提供大字版本、音频版本和其它语言版本，请拨打电话：0800 073 0510。

Ulotka dostępna jest również w dużym druku, wersji audio lub w innym języku. W tym celu zadzwoń pod numer 0800 073 0510.

ਇਹ ਪਰਚਾ ਵੱਡੇ ਅੱਖਰਾਂ, ਬੋਲ ਕੇ ਰੀਕਾਰਡ ਕੀਤਾ ਹੋਇਆ ਅਤੇ ਦੂਸਰੀਆਂ ਭਾਸ਼ਾਵਾਂ ਵਿਚ ਵੀ ਪ੍ਰਾਪਤ ਹੋ ਸਕਦਾ ਹੈ, 0800 073 0510 ਤੇ ਫੋਨ ਕਰੋ ਜੀ।

Aceasta broșura poate fi pusă la dispoziție tipărită cu caractere mari, versiune audio sau în alte limbi, pentru acest lucru va rugăm sunați la 0800 073 0510.

یہ کتابچہ آپ کو بڑے حروف کی لکھائی، سمعی صورت اور دیگر زبانوں میں مہیا کیا جا سکتا ہے۔ برائے مہربانی فون نمبر 08000730510 پر رابطہ کریں۔