

Radiological inserted gastrostomy (RIG)

Radiology Department Patient Information Leaflet

Introduction

This leaflet tells you about having a radiological inserted gastrostomy (RIG). It explains what is involved and what the possible risks are. It is not meant to replace informed discussion between you and your doctor, but can act as a starting point for such discussions. If you have any questions about the procedure, please ask the doctor who has referred you or the department which is going to perform it.

What is a gastrostomy?

A gastrostomy is a technique where a narrow plastic tube is placed through the skin into your stomach. Once in place, the tube can be used to give you liquid food directly into your stomach to provide nutrition. Gastrostomy tubes can be placed endoscopically (percutaneous endoscopic gastrostomy) or percutaneously (RIG). RIGs are placed by specially trained doctors called interventional radiologists.

Why do you need a gastrostomy?

You may be unable to eat or drink sufficient amounts to satisfy your nutritional needs, or there may be a problem with swallowing that makes it unsafe for you to eat or drink. There are several reasons for this.

You may have had a small plastic tube inserted through your nose, down into your stomach, to help with your feeding. This can only be left in place for a relatively short period of time. Obviously, if you do not receive enough nutrition, you will become very ill.

Are there any risks involved?

RIG is a very safe procedure, but as with any medical procedure, there are some risks and complications that can arise. Occasionally, it is not possible to place the tube into your stomach. This may require a different method of placement or, occasionally, you may need an operation to place the tube.

Sometimes, there is a leak around the tube. This is less likely to happen if the stomach has been attached to the muscles beneath the skin, but it can still sometimes occur. This can lead to the skin around the tube becoming very red, sore and painful (localised peritonitis). An attempt will be made to treat this, but it may become necessary to remove the tube to allow healing to occur, or an operation may be needed to sew up the hole in your stomach.

Very rarely, a blood vessel can be punctured accidentally when passing the needle into the stomach. This can result in bleeding. This may stop by itself, or if not, you may need a blood transfusion. Occasionally, it may require another procedure to block the bleeding artery or an operation to stop the bleeding. However, this is extremely rare.

Rarely, a bowel injury can occur as the stomach is very close to other loops of bowel. This may be treated conservatively, but sometimes bowel injury or leakage around the tube may need an operation and may cause a risk to life.

Who has made the decision?

The consultant in charge of your care and the interventional radiologist performing the procedure have discussed your case and feel that this is the best option.

However, you will also have the opportunity for your opinion to be considered and if, after discussion with your doctors, you no longer want the procedure, you can decide against it.

Are you required to make any special preparations?

A RIG is usually carried out as a day case procedure under local anaesthetic. You may be asked not to eat for four hours before the procedure, although you may still drink clear fluids, such as water. You may be asked to take a special shower and have antibiotics beforehand to reduce the chances of infection. If you have any allergies or have previously had a reaction to the dye (contrast agent), you **must** tell the radiology staff before you have the test.

Who will you see?

You will be seen by a specially trained team led by an interventional radiologist within the radiology department. Interventional radiologists have special expertise in reading the images and using imaging to guide catheters and wires to aid diagnosis and treatment.

Where will the procedure take place?

The procedure will take place in the interventional theatre located within the radiology department. This is similar to an operating theatre, into which specialised X-ray equipment has been installed.

What happens during the procedure?

You will be asked to get undressed and put on a hospital gown. A small cannula (thin tube) may be placed into a vein in your arm. You will lie flat on your back. You will have monitoring devices attached to your chest and finger, and you may be given oxygen. If you do not already have one inserted, a small tube will be placed through your nose into your stomach.

The procedure is performed under sterile conditions, and the interventional radiologist will wear a sterile gown and gloves to carry out the procedure. The skin below your ribs will be swabbed with antiseptic, and you will be covered with sterile drapes.

The interventional radiologist will use an ultrasound probe and X-rays once your stomach is filled with air to decide the best site for the RIG. Local anaesthetic will be injected into the skin to numb the area, and you will probably be given sedation. A small needle is placed into the stomach, through which a special feeding tube is placed. There are many different types of feeding tube available – some initially need stitches to keep them in place, others do not. Your interventional radiologist will discuss this with you.

Will it hurt?

When the local anaesthetic is injected, it will sting for a short while, but this soon wears off. There may be a little discomfort during the procedure, but any pain that you have will be controlled with painkillers. You may be aware of the tubes being passed into your stomach, but this should just be a feeling of pressure and not of pain.

How long will it take?

Every patient is different, and it is not always easy to predict. However, expect to be in the radiology department for about an hour.

What happens afterwards?

You will be taken back to your ward. Nursing staff will carry out routine observations including your pulse and blood pressure, and they will also check the treatment site. You will stay in bed for a couple of hours. The tube in your nose can be removed and the RIG can generally be used after a few hours. Your stomach may feel a little sore for a few days. If necessary, this can be controlled with painkillers.

How long will the tube stay in place?

This is a question that can only be answered by the doctors looking after you. It all depends on why you need the tube in the first place. You will need to discuss this with your consultant. The tube needs to stay in place until you can eat and drink safely and normally. In some cases, this might not be for a very long time.

You will have a specially trained dietitian looking after you who will show you how to look after the tube properly. If your tube has stitches holding it in place, these will be removed after about ten days. The tube should stay in by itself, although it is best covered with a light dressing, which the nurse looking after you can apply. Other tubes are kept in place by a small balloon. The nurse or dietitian caring for you will show you how to care for this balloon.

X-ray precautions

The procedure uses X-rays, and these use a small amount of radiation which may add slightly to the normal risk of cancer. Your doctor thinks that the benefit of the examination outweighs the risks. However, if you are concerned about these possible risks, please discuss these with this doctor.

For more information visit:

<https://www.gov.uk/government/publications/medical-radiation-patient-doses/patient-dose-information-guidance>

If you are a female patient, you must tell us if you are or might be pregnant before you attend for your scan.

Finally

Some of your questions should have been answered by this leaflet, but remember that this is only a starting point for discussion about your treatment with the doctors looking after you. Make sure you are satisfied that you have received enough information about the procedure.

Contact

The British Society of Interventional Radiology

www.bsir.org

Department of Radiology
Russells Hall Hospital
West Wing
Ground Floor
Pensnett Road
Dudley
West Midlands
DY1 2HQ

© This leaflet has been reproduced with kind permission from British Society of Interventional Radiology

If you have any questions, or if there is anything you do not understand about this leaflet, please contact:

The Radiology Department

Russells Hall Hospital switchboard number: 01384 456111

This leaflet can be downloaded or printed from:

<http://dgft.nhs.uk/services-and-wards/radiology>

If you have any feedback on this patient information leaflet, please email dgft.patient.information@nhs.net

This leaflet can be made available in large print, audio version and in other languages, please call 0800 073 0510.

للحصول على هذه النشرة بحجم أكبر، وعلى شكل إصدار صوتي و بلغات أخرى، الرجاء الاتصال بالرقم 08000730510.

此宣传单可提供大字版本、音频版本和其它语言版本，请拨打电话：0800 073 0510。

Ulotka dostępna jest również w dużym druku, wersji audio lub w innym języku. W tym celu zadzwoń pod numer 0800 073 0510.

ਇਹ ਪਰਚਾ ਵੱਡੇ ਅੱਖਰਾਂ، ਬੋਲ ਕੇ ਰੀਕਾਰਡ ਕੀਤਾ ਹੋਇਆ ਅਤੇ ਦੂਸਰੀਆਂ ਭਾਸ਼ਾਵਾਂ ਵਿਚ ਵੀ ਪ੍ਰਾਪਤ ਹੋ ਸਕਦਾ ਹੈ, 0800 073 0510 ਤੇ ਫੋਨ ਕਰੋ ਜੀ।

Aceasta broșura poate fi pusă la dispoziție tipărită cu caractere mari, versiune audio sau în alte limbi, pentru acest lucru va rugăm sunați la 0800 073 0510.

یہ کتابچہ آپ کو بڑے حروف کی لکھائی، سمعی صورت اور دیگر زبانوں میں مہیا کیا جا سکتا ہے۔ برائے مہربانی فون نمبر 08000730510 پر رابطہ کریں۔